



Rhythm Pharmaceuticals Announces New Employment Inducement Grants

December 7, 2022

BOSTON, Dec. 07, 2022 (GLOBE NEWSWIRE) -- Rhythm Pharmaceuticals, Inc. (Nasdaq: RYTM), a commercial-stage biopharmaceutical company focused on transforming the lives of patients and their families living with hyperphagia and severe obesity caused by rare melanocortin-4 receptor (MC4R) pathway diseases, today announced that on December 2, 2022, the Compensation Committee of Rhythm's board of directors granted inducement equity grants covering an aggregate of 29,925 shares of its common stock to three new employees, consisting of inducement stock options to purchase an aggregate of 19,950 shares of common stock and inducement restricted stock units, or RSUs, covering an aggregate of 9,975 shares of its common stock.

These inducement stock options and inducement RSUs are subject to the terms of the Rhythm Pharmaceuticals, Inc. 2022 Employment Inducement Plan (the "Inducement Plan").

The Inducement Plan is used exclusively for the grant of equity awards to individuals as an inducement material to the employees entering into employment with Rhythm pursuant to Nasdaq Listing Rule 5635(c)(4). The Inducement Plan was adopted by Rhythm's board of directors on February 9, 2022.

The stock options have an exercise price of \$ \$28.32 per share. Each option will vest as to 25% of the shares underlying such option on the first anniversary of the applicable date of hire of each individual, with the remaining 75% vesting in 12 equal quarterly installments over the three years thereafter, subject to each such employee's continued employment on each vesting date. The RSUs vest over four years, with 25% of the shares vesting on each anniversary of the applicable date of hire, subject to each such employee's continued employment on each vesting date.

About Rhythm Pharmaceuticals

Rhythm is a commercial-stage biopharmaceutical company committed to transforming the lives of patients and their families living with hyperphagia and severe obesity caused by rare melanocortin-4 receptor (MC4R) pathway diseases. Rhythm's precision medicine, setmelanotide, is approved by the U.S. Food and Drug Administration (FDA) for chronic weight management in adult and pediatric patients 6 years of age and older with monogenic or syndromic obesity due to pro-opiomelanocortin (POMC), proprotein convertase subtilisin/kexin type 1 (PCSK1) or leptin receptor (LEPR) deficiency confirmed by genetic testing, or patients with a clinical diagnosis of Bardet-Biedl syndrome (BBS). The European Commission (EC) and Great Britain's Medicines & Healthcare Products Regulatory Agency (MHRA) have authorized setmelanotide for the treatment of obesity and the control of hunger associated with genetically confirmed BBS or genetically confirmed loss-of-function biallelic POMC, including PCSK1, deficiency or biallelic LEPR deficiency in adults and children 6 years of age and above. Additionally, Rhythm is advancing a broad clinical development program for setmelanotide in other rare genetic diseases of obesity and is leveraging the Rhythm Engine and the largest known obesity DNA database -- now with approximately 45,000 sequencing samples -- to improve the understanding, diagnosis and care of people living with severe obesity due to certain genetic deficiencies. Rhythm's headquarters is in Boston, MA.

Setmelanotide Indication

In the United States, setmelanotide is indicated for chronic weight management in adult and pediatric patients 6 years of age and older with monogenic or syndromic obesity due to POMC, PCSK1 or LEPR deficiency as determined by an FDA-approved test demonstrating variants in POMC, PCSK1 or LEPR genes that are interpreted as pathogenic, likely pathogenic, or of uncertain significance (VUS) or BBS.

In the European Union, setmelanotide is indicated for the treatment of obesity and the control of hunger associated with genetically confirmed Bardet-Biedl syndrome (BBS) or genetically confirmed loss-of-function biallelic pro-opiomelanocortin (POMC), including PCSK1, deficiency or biallelic leptin receptor (LEPR) deficiency in adults and children 6 years of age and above.

Limitations of Use

In the United States and Europe, Setmelanotide should be prescribed and supervised by a physician with expertise in obesity with underlying genetic etiology.

Setmelanotide is not indicated for the treatment of patients with the following conditions as setmelanotide would not be expected to be effective:

- Obesity due to suspected POMC, PCSK1 or LEPR deficiency with *POMC*, *PCSK1* or *LEPR* variants classified as benign or likely benign
- Other types of obesity not related to POMC, PCSK1 or LEPR deficiency, or BBS, including obesity associated with other genetic syndromes and general (polygenic) obesity.

WARNINGS AND PRECAUTIONS

Skin Monitoring: Setmelanotide may lead to generalized increased skin pigmentation and darkening of pre-existing naevi because of its pharmacologic effect. Full body skin examinations should be conducted annually to monitor pre-existing and new skin pigmentary lesions before and during treatment with setmelanotide.

Heart rate and blood pressure monitoring: Heart rate and blood pressure should be monitored as part of standard clinical practice at each medical visit (at least every 6 months) for patients treated with setmelanotide.

Prolonged penile erection: Spontaneous penile erections have been reported in clinical trials with setmelanotide. Patients who have a penile erection lasting longer than 4 hours should be instructed to seek emergency medical attention for potential treatment of priapism.

Depression: In clinical trials, depression has been reported in patients treated with setmelanotide. Patients with depression should be monitored at each medical visit during treatment with setmelanotide. Consideration should be given to discontinuing setmelanotide if patients experience suicidal thoughts or behaviors.

Pediatric Population: The prescribing physician should periodically assess response to setmelanotide therapy. In growing children, the impact of weight loss on growth and maturation should be evaluated. The prescribing physician should monitor growth (height and weight) using age- and sex-appropriate growth curves.

Excipients: This medicinal product contains 10 mg benzyl alcohol in each ml. Benzyl alcohol may cause allergic reactions. Patients who are pregnant or breastfeeding should be advised of the potential risk from the excipient benzyl alcohol, which might accumulate over time and cause metabolic acidosis. This medicinal product should be used with caution in patients with hepatic or renal impairment, because of the potential risk from the excipient benzyl alcohol which might accumulate over time and cause metabolic acidosis.

Sodium: This medicinal product contains less than 1 mmol sodium (23 mg) per dose, that is to say essentially “sodium-free.”

ADVERSE REACTIONS

The most frequent adverse reactions are hyperpigmentation (51%), injection site reaction (39%), nausea (33%), and headache (26%).

USE IN SPECIFIC POPULATIONS

Pregnancy

There are no data from the use of setmelanotide in pregnant women. Animal studies do not indicate direct harmful effects with respect to reproductive toxicity. However, administration of setmelanotide to pregnant rabbits resulted in decreased maternal food consumption leading to embryo-foetal effects. As a precautionary measure, setmelanotide should not be started during pregnancy or while attempting to get pregnant as weight loss during pregnancy may result in fetal harm. If a patient who is taking setmelanotide has reached a stable weight and becomes pregnant, consideration should be given to maintaining setmelanotide treatment as there was no proof of teratogenicity in the nonclinical data. If a patient who is taking setmelanotide and still losing weight gets pregnant, setmelanotide should either be discontinued, or the dose reduced while monitoring for the recommended weight gain during pregnancy. The treating physician should carefully monitor weight during pregnancy in a patient taking setmelanotide.

Breast-feeding

It is unknown whether setmelanotide is excreted in human milk. A nonclinical study showed that setmelanotide is excreted in the milk of nursing rats. No quantifiable setmelanotide concentrations were detected in plasma from nursing pups. A risk to the newborn/infant cannot be excluded. A decision must be made whether to discontinue breastfeeding or to discontinue/abstain from setmelanotide therapy taking into account the benefit of breastfeeding for the child and the benefit of therapy for the mother.

Fertility

No human data on the effect of setmelanotide on fertility are available. Animal studies did not indicate harmful effects with respect to fertility.

To report SUSPECTED ADVERSE REACTIONS, contact Rhythm Pharmaceuticals at +1 (833) 789-6337. See [Summary of Product Characteristics' APPENDIX V](#) for a list of European national reporting systems to communicate adverse reactions.

Please see the full Prescribing Information for additional Important Safety Information.

Forward-Looking Statements

This press release contains forward-looking statements within the meaning of the Private Securities Litigation Reform Act of 1995. All statements contained in this press release that do not relate to matters of historical fact should be considered forward-looking statements, including without limitation statements regarding the potential, safety, efficacy, and regulatory and clinical progress, potential regulatory submissions, approvals and timing thereof of setmelanotide, and our business strategy and plans, including regarding commercialization of IMCIVREE. Statements using word such as “expect”, “anticipate”, “believe”, “may”, “will” and similar terms are also forward-looking statements. Such statements are subject to numerous risks and uncertainties, including, but not limited to, our ability to enroll patients in clinical trials, the design and outcome of clinical trials, the impact of competition, the ability to achieve or obtain necessary regulatory approvals, risks associated with data analysis and reporting, our ability to successfully commercialize setmelanotide, our liquidity and expenses, the impact of the COVID-19 pandemic on our business and operations, including our preclinical studies, clinical trials and commercialization prospects, and general economic conditions, and the other important factors discussed under the caption “Risk Factors” in our Quarterly Report on Form 10-Q for the three months ended September 30, 2022 and our other filings with the Securities and Exchange Commission. Except as required by law, we undertake no obligations to make any revisions to the forward-looking statements contained in this release or to update them to reflect events or circumstances occurring after the date of this release, whether as a result of new information, future developments or otherwise.

Corporate Contact:

David Connolly
Head of Investor Relations and Corporate Communications
Rhythm Pharmaceuticals, Inc.
857-264-4280
dconnolly@rhythmtx.com

Investor Contact:

Hannah Deresiewicz
Stern Investor Relations, Inc.
212-362-1200
hannah.deresiewicz@sternir.com

Media Contact:

Adam Daley
Berry & Company Public Relations
212-253-8881
adaley@berrypr.com



Source: Rhythm Pharmaceuticals, Inc.